

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 July 2004 (29.07.2004)

PCT

(10) International Publication Number
WO 2004/063149 A1

(51) International Patent Classification⁷: **C07C 317/44**,
315/02, 323/52, 319/14

(21) International Application Number:
PCT/IB2004/000002

(22) International Filing Date: 8 January 2004 (08.01.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03290082.1 13 January 2003 (13.01.2003) EP

(71) Applicant (for all designated States except US): **ORGANISATION DE SYNTHÈSE MONDIALE ORSY-MONDE** [FR/FR]; 5 rue Charles Martigny, F-94700 MAISONS ALFORT (FR).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ROSE, Sébastien** [FR/FR]; 23, chemin du Tour de Ville, F-60190 ARSY (FR). **KLEIN, Dominique** [FR/FR]; 28, rue Itasse, F-78780 MAURECOURT (FR).

(74) Agents: **BERNASCONI, Jean** et al.; Cabinet LAVOIX, 2, Place d'Estienne d'Orves, F-75441 PARIS (FR).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR PREPARING METHYL 2-DIPHENYLMETHYLSULFINYLACETATE

(57) Abstract: Method for preparing methyl 2-diphenylmethylsulfinylacetate (MDMSA) comprising the steps of (i) conversion of benzhydrol into methyl 2-diphenylmethylthioacetate; and (ii) conversion of methyl 2-diphenylmethylthioacetate into methyl 2-diphenylmethylsulfinylacetate by oxidation.

THIS PAGE BLANK (USPTO)

METHOD FOR PREPARING METHYL 2-DIPHENYLMETHYLSULFINYLACETATE

The present invention relates to a novel method for preparing methyl 2-diphenylmethylsulfinylacetate (MDMSA).

MDMSA is disclosed as an intermediate compound in the synthesis of modafinil also known as 2-[(diphenylmethyl)sulfinyl]acetamide. Modafinil which is a synthetic acetamide with wake-promoting activity, is useful in the treatment of narcolepsy, among other disorders.

The inventors have now discovered a novel route for synthesizing MDMSA which is applicable at the industrial scale.

Advantageously, the MDMSA can be obtained in two or three steps, each being characterized by high yields.


In an advantageous embodiment, these steps may be carried out in the same reactor and the same solvent, without isolating the intermediate compounds.

The aim of the present invention is to provide an economical and efficient method for preparing MDMSA.

These aims and others are achieved by the present invention which relates to a method for preparing methyl 2-diphenylmethylsulfinylacetate comprising the steps of:

- (i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and
- (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate by oxidation.

Scheme 1 illustrates in general the steps used in this method :



Scheme 1

The reactions of steps (i) and (ii) used in this method are carried out in an appropriate solvent which may be easily chosen by persons skilled in the art, it being understood that an appropriate solvent denotes a solvent which is nonreactive towards the starting reagents, the intermediates or the products, at the reaction temperature considered, it being possible for the latter to vary from the solidification point of the solvent to the boiling point of the solvent.

A given reaction may be carried out in a solvent or in a mixture of several solvents, the solvent(s) being generally chosen according to the type of reaction considered and the subsequent treatment of the reaction medium.

In a preferred embodiment, the solvent is an aprotic solvent.

By way of illustration and without limitation of aprotic solvents which may be suitable for the method according to the invention, there may be mentioned in particular chlorinated solvents, aromatic solvents, hydrocarbon solvents and ethereal solvents.

Among the chlorinated solvents, chloroform, dichloromethane or chlorobenzene may be mentioned in particular.

Among the appropriate aromatic solvents, there may be mentioned, for example, benzene, toluene and chlorobenzene.

As examples of appropriate hydrocarbon solvents, cyclohexane, pentane and hexane may be mentioned.

Solvents such as diethyl ether, tetrahydrofuran and dioxane are useful as ethereal solvents.

Step (i) and (ii) may be conducted separately, notably in different solvents, where each intermediate is independently isolated.

In an advantageous embodiment, the reaction steps are conducted in the same reactor and without isolation of any intermediates.


STEP i)

In a preferred embodiment, the conversion of benzhydrol into methyl diphenylmethylthioacetate comprises two steps of :

a1) conversion of benzhydrol into a benzhydryl carboxylate ; and

b1) conversion of benzhydryl carboxylate into methyl diphenylmethyl thioacetate.

Scheme 2 illustrates in general steps a1) and b1) used in this method :



Scheme 2

Step a1)

In one preferred embodiment, step a1) comprises reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate solvent, preferably aprotic.

In the context of the present invention, dichloromethane is particularly preferred for its advantageous properties of extraction, distillation at low temperature, its nonflammability, its chemical neutrality and its capacity to be easily recycled in the context of this method.

Generally, the acid anhydride and the aprotic solvent are loaded simultaneously into the reactor, at a temperature of about 20°C.

The acid anhydrides for the purposes of the present description denote either symmetric anhydrides of monocarboxylic acids, mixed anhydrides, or cyclic anhydrides of polycarboxylic acids.

By way of examples of acid anhydrides which may be suitable in step a1) according to the invention, there may be mentioned in particular acetic anhydride, butyric anhydride and propanoic anhydride, acetic anhydride being particularly preferred.

The preferred reaction conditions are those which provide for the use of an equimolar quantity of acid anhydride relative to the benzhydrol in the aprotic solvent.

However, it is preferable to carry out the procedure in the presence of a slight excess of acid anhydride. An excessively high excess may indeed induce the formation of by-products in the subsequent steps of the method while an excessively small quantity is capable of slowing the kinetics of the oxidation reaction carried out in step (ii). Thus, the molar ratio of acid anhydride preferably varies between 1 and 1.2, even better between 1 and 1.1, and is advantageously

1.05, which corresponds to the optimum molar ratio of acid anhydride which makes it possible to obtain a clean synthesis.

Preferably, the procedure is carried out in the presence of a volume of solvent varying from 1.5 to 5 volumes, optimally in the presence of about 2
5 volumes. Under these conditions, the dilution brought about promotes the formation of the carboxylate.

According to an equally preferred variant of step a), the inorganic acid used is chosen from hydrochloric acid, hydrobromic acid, o-phosphoric acid and sulphuric acid, sulphuric acid being particularly preferred, in particular in the form
10 of a 96% aqueous solution.

Preferably, the procedure is carried out in the presence of a quantity of inorganic acid ranging from 0.02 to 0.3 molar equivalents relative to the benzhydrol, even better from 0.05 to 0.15 equivalents.

As a safety precaution, the inorganic acid is generally introduced at a
15 temperature of about 0°C so as to control the exothermicity of the reaction.

The benzhydrol is then loaded at a sufficient temperature to allow a sufficiently rapid reaction kinetics, but not too high so as to avoid the formation of by-products such as benzhydryl ether. It is particularly preferable to carry out the procedure at a temperature of between -5°C and +5°C, even better between -2°C
20 and +2°C for an introduction time ranging from 45 minutes to 2 hours, preferably in about 1 hour. This introduction time indeed makes it possible to control the exothermicity of the reaction and to limit the formation of by-products.

The benzhydrol is generally maintained at this temperature for a sufficient contact time to obtain a complete reaction, but not too high in order to avoid the degradation of the benzhydryl carboxylate. The expression "complete reaction",
25 for the purposes of the present invention, is understood to mean a reaction leading to the production of the derived product with a conversion rate greater than 99.2%, and preferably greater than 99.5%. In general, a complete reaction is obtained after a contact time of 2 hours.

30 The benzhydryl carboxylate obtained can be immediately used in the next step, without intermediate isolation.

Step a1) can also be realized by any other appropriate method.

As an example, step a1) can be realized by reacting benzhydrol with a

carboxylic acid, for instance :

- 2-methyl-butiric acid in combination with :
 - o SOCl_2 , pyridine, in benzene ; or
 - o H_2SO_4 in dichloromethane ; or
 - o TsOH in benzene

as disclosed in the reference : Fujita S. and al., Bull. Chem. Soc. Jpn 1972, 45 : 2571-2574 ;

- or acetic acid in combination with potassium iodide as described in Strazzoli P. et al., Recl. Trav. Chim., The Netherlands, 1991 ; 1:5-12.

By way of example, step a1) can also be realized by reacting a carboxylic acid salt, for instance acetic acid sodium salt as disclosed in Herzig S., Justus Liebigs, Ann. Chem. ; 1921 ; 422:330.

By way of example, step a1) can also be realized by reacting benzhydrol with a carboxylic acid chloride, as for example acetyl chloride in the presence of triethylamine as disclosed in Roizel B. and al., Chem. Commun., 2000, 16:1507-1508.

As an example, step a1) can also be realized by reacting benzhydrol with acetic acid ethyl ester in the presence of $\text{Ti}(\text{OC}_2\text{H}_5)_4$ (Schnurrenberger P. and al., Helv. Chim. Acta, 1982, 65(4) : 1197-1201) ; or of iron (III) perchlorate (ITP) (Kumar B. and al., Indian J. Chem. Sect. B, 1993, 32(2) : 292-293) ; or of $\text{Fe}(\text{ClO}_4)_3$, SiO_2 (Parmar and al., Synth. Commun., 1999, 29(1) : 139-144).

Step b1)

Step b1) can be performed by any appropriate method.

In a preferred embodiment, step b1) comprises bringing the solution obtained in step a1) into contact with methyl thioglycolate.

The methyl thioglycolate is generally introduced in about 10 minutes at 0°C but with no temperature restriction (it can then increase to about 9°C). The reaction medium is then heated to a sufficiently high temperature to drive the reaction kinetics, but not too high so as to avoid the formation of by-products. Generally, the procedure is carried out at a temperature of between 15°C and 25°C , preferably between 18°C and 22°C , and contact is maintained at this

temperature for a sufficient time to obtain complete reaction, with few by-products, generally for 2 to 3 hours, preferably for a period of 2 hours.

The methyl diphenylthioacetate can be used in step (ii) without intermediate isolation.

5


In another preferred embodiment, step i) comprises two steps of :

a2) conversion of benzhydrol into benzhydryl carboxylic acid ;

b2) conversion of benzhydryl carboxylic acid into methyl diphenylmethylthioacetate.

10

This method is illustrated by scheme 3 :



Scheme 3

15

Step a2)

Step a2) may be realized according to any appropriate method and notably according to conditions disclosed in : Dahlbom O., Acta Chem. Scand., 1948, 2 : 856-858 ; Carceller E. et al., J. Med. Chem., 1993; 36 : 2984-2997 ; Lisac S. et al., J. Organomet. Chem . 1996, 507 : 215-220 ; Okarvi S. et al., J. Labelled Compd. Radiopharm, 1997, 39 : 853-874 ; Patent Thomae GmbH DE 2812542, 1979, Chem. Abstract 1980 ; 92 ; 198165 ; Iskander, Y. et al., J. Chem. Soc., 1961, 2397-2402.

In a particular embodiment, step a2) is realized by reacting benzhydrol with thioacetic acid, in the presence of an organic or inorganic acid.

Preferably, the solvent is a protic solvent, more preferably a carboxylic acid and notably acetic acid.

Inorganic or organic acid is preferably chosen among hydrochloric acid, POCl₃, trifluoroacetic acid, hydrobromic acid, o-phosphoric acid, sulphuric acid,

POCl_3 and trifluoroacetic acid being particularly preferred.

Preferably, the reaction is realized at room temperature.


Step b2)

5 Esterification reaction of step b2) may be realized by any methods known from the person skilled in the art.

In another particular embodiment, step (i) comprises two steps of :

- 10 a3) conversion of the hydroxyl group of benzhydrol into a leaving group ;
b3) conversion of the obtained product into methyldiphenyl-
methylthioacetate.

This method is illustrated by scheme 4 :



15

Scheme 4

Step a3)

The leaving group in step a3) means any group that can be removed easily by a nucleophilic reactant. Leaving groups may be selected from the group consisting of halogenes, such as chloro- and bromo- radicals, or sulfonyl groups
20 such as methanesulfonyl - or *p*-toluenesulfonyl- radicals.

Step a3) may be realized by any methods known from the person skilled in the art.

25 As an example, hydroxyl group of benzhydrol may be converted into chloro- or bromo- radical by reacting benzhydrol with thionyl chloride or thionyl bromide.


As an example, hydroxyl group of benzhydrol may be converted into methanesulfonate group or into *p*-toluenesulfonate group by reacting benzhydrol respectively with methanesulfonyl chloride or *p*-toluenesulfonyl chloride.

Step b3)

In a preferred embodiment, step b3) is realized according to conditions of step b1).

5 Step b3) may also be realized by any other appropriate method.

In another preferred embodiment, step i) comprises reacting benzhydrol with methylthioglycolate in the presence of a metallic catalyst in a solvent, as illustrated by scheme 5 :



10

Scheme 5

Preferably, the metallic catalyst is chosen from ZnCl₂, ZnBr₂, ZnI₂, ZnI₂ being particularly preferred.

Preferably, the solvent is chosen from aprotic solvents, more preferably from halogenated solvent and particularly chlorinated solvents such as dichloromethane, dichloroethane.

15

STEP ii)

Inventors have now settled oxidation conditions that allow interestingly to control the conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate, notably formation of by-products, yields and kinetics of the reaction.

20

Various parameters used in step (ii) capable of influencing the efficiency of the reaction may be optimized, such as the quantity of oxidizing agent introduced, the dilution of the reaction medium, the reaction temperature, the contact time and the acidity of the reaction medium.

25

Thus, in general, an excessively small quantity of oxidizing agent means an incomplete reaction. Conversely, an excessively large excess promotes the

formation of the dioxidized by-product, sulfone.

By way of illustration of oxidizing agents which may be suitable for the invention, there may be mentioned in particular oxone, potassium permanganate, sodium percarbonate, peroxides such as hydrogen peroxide, tert-butyl hydroperoxide and m-chloroperoxybenzoic acid, hydrogen peroxide being particularly preferred.

In a preferred embodiment, step ii) is performed in dichloromethane.

According to a preferred embodiment, the oxidizing agent used in step (ii) is hydrogen peroxide, preferably in the form of a 35% aqueous solution. Indeed, a lower titre causes a higher dilution, which can decrease the kinetics of the reaction.

Although a stoichiometric quantity of oxidizing agent is sufficient, it is preferably to carry out the procedure in the presence of a slight excess, preferably in the presence of a molar ratio of between 1 and 1.1.

The acidity of the medium results from the operating conditions of step (i).

The reaction temperature can influence the kinetics of the oxidation reaction. Thus, it is preferable that the temperature is between 28°C and 37°C, the temperature range for which the reaction kinetics is particularly increased taking into account the acidity of the medium.

Preferably, a reaction temperature of between 28°C and 32°C is preferred. This temperature indeed makes it possible to have optimum control of the method, in particular the stopping point beyond which superoxidation becomes non-negligible.

The contact time in order to obtain a clean and complete reaction may vary according to the scale of operation and also according to the quantity of inorganic acid, in particular of sulfuric acid, present in the reaction medium at step b).

Preferably, the reaction is considered as being "complete" in step (ii) when the ratio $R_1 = \text{nonoxidized derivative} / (\text{monooxidized} + \text{dioxidized} + \text{nonoxidized derivative}) < 0.5\%$.

The expression "clean" reaction is understood to mean, for the purposes of the present description, a reaction in which the ratio $R_2 = \text{dioxidized derivative} / (\text{monooxidized} + \text{dioxidized} + \text{nonoxidized derivative}) < 0.5\%$.

The contact time necessary in order to obtain a clean and complete

reaction may be determined using conventional analytical techniques which make it possible to monitor the progress of the reaction, such as HPLC (High-Performance Liquid Chromatography), IR (Infrared) or NMR (Nuclear Magnetic Resonance).

5 In general, the contact time necessary and sufficient to obtain a clean and complete reaction is less than 35 hours, preferably less than or equal to 33 hours, so as to avoid the formation of by-products such as in particular diphenylmethylthioacetic acid, and greater than 20 hours, preferably greater than or equal to 25 hours.

10 Of course, it is within the capability of persons skilled in the art to adjust the contact time necessary for obtaining a complete reaction at the scale of operation considered.

An increase in the quantity of inorganic acid in the reaction medium can nevertheless make it possible to significantly reduce the oxidation reaction time.

15 Without wishing to be limited to any theory, a hypothesis which makes it possible to explain this unexpected effect is that the inorganic acid plays a catalyst role in the mechanism of oxidation by the oxidizing agent. By way of example, in the case of sulphuric acid, it is assumed that a reaction intermediate of the H_2SO_5 type is formed in the medium, transferring oxygen either directly to
20 the oxidizable species, or indirectly by an accelerated formation of peracetic acid.

Thus, according to a preferred variant of step (ii) of the method according to the invention, an additional quantity of inorganic acid, preferably of 0.02 to 0.3 molar equivalents, and more preferably of 0.05 to 0.15 molar equivalents is added to the reaction medium in step b1), generally prior to the introduction of the
25 oxidizing agent. An acceleration of the reaction kinetics is then observed.

Advantageously, the contact time required which is sufficient to obtain a complete and clean reaction in step (ii) is considerably reduced and is generally between 10 and 13 hours.

Advantageously, the introduction of the inorganic acid in two portions
30 makes it possible to reduce the acidity of the reaction medium in steps a1) and b1) and therefore to limit the formation of by-products.

STEP (iii)

In a further embodiment, the method according to the invention comprises an additional step of iii) recovering the methyl 2-diphenyl-methylsulfinylacetate obtained.

5

The isolation of the MDMSA formed in step (iii) can be carried out according to any conventional method known to persons skilled in the art.

Preferably, the MDMSA is isolated by extraction.

The organic phases are then combined and concentrated under reduced
10 pressure, preferably at a temperature of 70°C.

According to a particular variant, the solvent is distilled to dryness.

The product may be purified according to any method known to persons skilled in the art such as recrystallization or chromatography.

According to a particular embodiment, step (iii) may comprise a step of
15 direct crystallization of the MDMSA.

The expression "direct crystallization", for the purposes of the present description, is understood to mean a crystallization of the noncrystallized product caused by the addition of appropriate solvents, preferably chosen in particular from methanol, ethanol, ethyl acetate, isopropyl acetate and toluene, isopropyl
20 acetate being particularly preferred.

In this context, the crystallization solvent is introduced after substantial removal of the aprotic solvent.

Advantageously, this direct crystallization makes it possible to purify the crude product immediately in the remainder of the method and thus to dispense
25 with an isolation step and a more costly subsequent retreatment step.

In a particularly preferred embodiment, the method comprises the steps of:

- i) a1) converting benzhydrol into a benzhydryl carboxylate by reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate aprotic solvent ;
30 b1) converting the benzhydryl carboxylate into methyl diphenylmethylthioacetate by bringing the above solution into contact with methyl thioglycolate;
- ii) converting the diphenylmethylthioacetate into methyl 2-


diphenylmethanethiolacetate by bringing the above solution into contact with an oxidizing agent ;

and optionally ;

iii) recovering the methyl 2-diphenylmethanethiolacetate obtained.

5

This embodiment is illustrated by scheme 6 :



Scheme 6

10 Advantageously, the MDMSA is obtained in three successive steps, each being characterized by high yields.

Furthermore, these three steps may be carried out in the same reactor and the same solvent, without isolating the intermediate compounds.

15 Advantageously, the formation of undesirable by-products is limited and controlled according to this method, which makes it possible to dispense with subsequent reprocessing steps.

Finally, according to another advantageous aspect, this method makes it possible to reduce the costs of manufacturing MDMSA, in particular because of its ease of use, and its high productivity and reproducibility.

20

The present invention also aims at the MDMSA obtained by the method of the invention and particularly by the method implementing :

i) a1) converting benzhydrol into a benzhydryl carboxylate by reacting benzhydrol and an acid anhydride in the presence of

an inorganic acid and in an appropriate aprotic solvent ;

b1) converting the benzhydryl carboxylate into methyl diphenylmethylthioacetate by bringing the above solution into contact with methyl thioglycolate;

5 ii) converting the diphenylmethylthioacetate into methyl 2-diphenylmethylsulfinylacetate by bringing the above solution into contact with an oxidizing agent ;

and optionally ;

10 iii) recovering the methyl 2-diphenylmethylsulfinylacetate obtained.

The present invention is also directed to a method for preparing modafinil comprising the steps of :

(i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and

15 (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate by oxidation.

EXAMPLES

Example 1. – Synthesis of MDMSA on the laboratory scale (0.5 l)

5 a) Benzhydryl acetate

108.3 g (1.05 mol; 1.05 eq) of acetic anhydride are diluted in 370 ml of dichloromethane at 20°C. The solution obtained is cooled to $0 \pm 2^\circ\text{C}$ before introducing 2.8 ml of a 96% sulphuric acid solution in a time of about 10 minutes. After stirring for about 10 minutes, 184.2 g (1 mol, 1 eq) of benzhydrol are
10 introduced in portions at $0^\circ\text{C} \pm 2^\circ\text{C}$ in 60 ± 15 minutes. The reaction medium is maintained in contact for 2 hours at this temperature.

b) Methyl diphenylmethylthioacetate (MDMTA)

108.3 g (1.02 mol; 1.02 eq) of methyl thioglycolate are introduced at $0^\circ\text{C} \pm 2^\circ\text{C}$, and then the reaction mixture is heated to $20^\circ\text{C} \pm 2^\circ\text{C}$ and kept in contact for
15 2 hours at this temperature.

c) Methyl 2-diphenylmethylsulfinylacetate (MDMSA)

The reaction medium is heated to $30^\circ\text{C} \pm 2^\circ\text{C}$, and 100.5 ± 0.5 g of a 35%
20 hydrogen peroxide solution are added. The progress of the oxidation reaction is monitored by HPLC. After 25 hours of contact, the reaction is complete (R_1 and $R_2 < 0.5\%$).

d) Recovery of MDMSA

25 150 ml of distilled water are stirred. A 25% aqueous ammonia solution is added so as to bring the pH to 8. The aqueous organic phases are separated, and the aqueous phase is extracted with 2 x 100 ml of dichloromethane. The organic phases are then combined and concentrated under reduced pressure at a temperature of 70°C . The concentrate obtained crystallizes to give, after
30 grinding, a white powder with a yield of $98.0 \pm 0.5\%$.

Examples 2 and 3. – Synthesis of MDMSA on a pilot scale (100 l)

Example 2: Synthesis of crystallized MDMSA

a) Benzhydryl acetate

5 A solution of acetic anhydride (8.73 kg; 85.5 mol; 1.05 eq) in methylene chloride (40 l) is prepared at $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$. A 96% sulphuric acid solution (225 ml; 4.1 mol; 0.05 eq) is then added with a stirring of 100 revolutions/min, at $0^{\circ}\text{C} \pm 5^{\circ}\text{C}$ in 5 to 10 minutes. The benzhydrol (15 kg; 81.4 mol, 1 eq) is then introduced at $0^{\circ}\text{C} \pm 2^{\circ}\text{C}$ in 1.25 hours. The reaction mixture is stirred for a contact time of 2
10 hours.

b) Methyl diphenylmethylthioacetate (MDMTA)

 Methyl thioglycolate (8.81 kg; 82.9 mol; 1.02 eq) is introduced at $0^{\circ}\text{C} \pm 2^{\circ}\text{C}$ in about 10 minutes. The reaction mixture is brought to the temperature of $20^{\circ}\text{C} \pm$
15 2°C and kept at this temperature for a contact time of 2 hours, with a stirring of 100 revolutions/min.

c) Methyl 2-diphenylmethylsulfinylacetate (MDMSA)

 The reaction mixture is then brought to 30°C before introducing sulphuric
20 acid (450 ml; 8.1 mol; 0.1 eq) with stirring (100 revolutions/min), in about 5 to 10 minutes. A 35% hydrogen peroxide solution (8.19 kg; 84.3 mol; 1.035 eq) is then introduced at $30^{\circ}\text{C} \pm 2^{\circ}\text{C}$ in 1 hour. The contact time is determined by monitoring the reaction by HPLC (cf. Table 1).

25 d) Recovery of MDMSA

 The mixture is cooled to $20^{\circ}\text{C} \pm 2^{\circ}\text{C}$ and then 20 l of water are introduced. After neutralizing the reaction medium by adding a sufficient quantity of NH_4OH so that $8 < \text{pH} < 9$, the aqueous and organic phases are separated and the aqueous phase extracted twice with 10 l of methylene chloride. The chlorinated
30 phases are washed with 10 l of water.

 The solvent is distilled under atmospheric pressure and then under reduced pressure, at a jacket temperature of 70°C . When the distillation is

complete, isopropyl acetate (1.8 vol; 42 l) is added and the whole is cooled to -10°C. After draining and drying under vacuum at 45°C, MDMSA is obtained.

Table 1: Yield and quality of the MDMSA and of the intermediate products
5 obtained by this method for trials 1 to 4 (step (ii)).

Trial	Formation of benzhydryl acetate		Formation of MDMA		Formation of MDMSA		HPLC of the finished product	Yield
	Time	HPLC %	Time	HPLC %	Time	HPLC %	%	
1	2h	98.9	2 h	98.5	10h	T = 1 S = 0.3	D = 99.7 S = 0.3	88
					11h	T = 0.7 S = 0.35		
					11h30	T = 0.3 S = 0.4		
2	2h	99.3	2h	99.1	11h30	T = 0.9 S = 0.35	D = 99.6 S = 0.3 T = 0.1	86.4
3	2h	99.3	2h	99.0	11h30	T = 0 S = 0.4	D = 99.7 S = 0.3	90.1
4	2h	99.3	2h	99.1	5 h	T = 7.2 S = 0.16	D = 99.7 S = 0.3	90.0
					6h30	T = 4.4 S = 0.17		
					8h	T = 2.3 S = 0.22		
					9h	T = 1.6 S = 0.14		
					10h	T = 1.15 S = 0.3		
					11h	T = 0.6 S = 0.38		
					11h30	T = 0.3 S = 0.38		

D = MDMSA

S = Sulphone

T = MDMA

10 These results demonstrate that the production of benzhydryl acetate and of MDMA can be reproducible.

The formation of MDMSA takes place in 11 h 30 min and reproducibly gives, after crystallization from isopropyl acetate, a final product which is in conformity (R_1 and $R_2 < 0.5\%$), with a yield of the order of 90%.

5 *Example 3: Synthesis of crude MDMSA*

a) Benzhydryl acetate

A solution of acetic anhydride (8.73 kg; 85.5 mol; 1.05 eq) in methylene chloride (40 l) is prepared at 20°C. A 96% sulphuric acid solution (225 ml; 4.1 mol; 0.05 eq) is then added with a stirring of 100 revolutions/min, at 0°C ± 5°C
10 in 5 to 10 minutes. The benzhydrol (15 kg; 8.4 mol, 1 eq) is then introduced at 0°C ± 2°C in 1.25 hours. The reaction mixture is stirred for a contact time of 2 hours.

b) Methyl diphenylmethylthioacetate (MDMTA)

15 Methyl thioglycolate (8.81 kg; 82.9 mol; 1.02 eq) is introduced at 0°C in about 10 minutes. The reaction mixture is brought to the temperature of 20°C ± 2°C and kept at this temperature for a contact time of 2 hours, with stirring of 100 revolutions/min.

20 c) Methyl 2-diphenylmethylsulfinylacetate (MDMSA)

Once the reaction mixture has been brought to 30°C ± 2°C, a 35% hydrogen peroxide solution (8.19 kg; 84.3 mol; 1.035 eq) is introduced with stirring (100 revolutions/min) in 1 hour. The contact time is determined by monitoring the reaction by HPLC (cf. Table 2).

25

d) Recovery of MDMSA

The mixture is cooled to 20°C and then 20 l of water are introduced. After neutralizing the reaction medium by adding a sufficient quantity of NH₄OH so that 8 < pH < 9, the aqueous and organic phases are separated and the aqueous
30 phase extracted twice with 10 l of methylene chloride. The chlorinated phases are washed with 10 l of water.

The solvent is distilled to dryness under atmospheric pressure and then under reduced pressure at a jacket temperature of 70°C in a Moritz® turbosphere.

Table 2: Yield and quality of the finished product and of the intermediate products for trials 5 and 6

Trial	Formation of benzhydryl acetate		Formation of MDMA		Formation of MDMSA		HPLC of the finished product	Yield
	Time	HPLC %	Time	HPLC %	Time	HPLC %	%	
5	2h	99.6	2h	99.2	35h	T = 0.3 S = 0.4	D = 99.12 S = 0.44 T = 0.24	97.2
6	2h	99.3	2h	98.9	33h	T = 0.17 S = 0.4	D = 99.2 S = 0.4 T = 0.2	97

5 D = MDMSA

S = Sulphone

T = MDMA

10 These results show that the steps of formation of benzhydryl acetate and of MDMA can be reproducible.

The step of oxidation of MDMA requires a contact time of about 33-35 hours and gives an MDMSA product which is in conformity (R_1 and $R_2 < 0.5\%$), with good yields (of the order of 97%).

CLAIMS

1. Method for preparing methyl 2-diphenylmethylsulfinylacetate (MDMSA) comprising the steps of :

- 5 (i) conversion of benzhydrol into methyldiphenylmethylthioacetate ; and
- (ii) conversion of methyldiphenylmethylthioacetate into methyl-2-diphenylmethylsulfinylacetate.

10 2. Method according to claim 1, in which step (i) comprises the following steps :

- a1) conversion of benzhydrol to benzhydrol carboxylate in an appropriate solvent ;
- b1) conversion of the benzhydrol carboxylate to methyl
- 15 diphenylmethylthioacetate.

3. Method according to claim 2, in which the step (a1) comprises reacting benzhydrol and an acid anhydride in the presence of an inorganic acid and in an appropriate solvent.

20

4. Method according to claim 3, in which the solvent is an aprotic solvent.

5. Method according to claim 4, in which the aprotic solvent is chosen from chlorinated solvents, aromatic solvents, hydrocarbon solvents and ethereal

25 solvents.

6. Method according to claim 5, in which the aprotic solvent is chosen from chlorinated solvents.

30 7. Method according to claim 6, in which the solvent is dichloromethane.

8. Method according to any one of claims 3 to 7, in which the acid anhydride is chosen from acetic anhydride, propanoic anhydride and butyric anhydride.

5 9. Method according to claim 8, in which the acid anhydride is acetic anhydride.

10 10. Method according to any one of claims 3 to 9, in which the inorganic acid is chosen from hydrochloric acid, butyric acid, o-phosphoric acid and sulfuric acid.

11. Method according to claim 10, in which the inorganic acid is sulfuric acid.

15 12. Method according to any one of claims 3 to 11, in which the quantity of inorganic acid used is from 0.02 to 0.3 molar equivalents relative to the benzhydrol.

20 13. Method according to any one of claims 3 to 12, in which the reaction temperature in step a) is between -5°C and $+5^{\circ}\text{C}$.

14. Method according to claim 2 to 13, in which step b1) comprises bringing the solution obtained in step a) into contact with methyl thioglycolate.

25 15. Method according to claim 14, in which the contact time used in step b1) is between 2 and 3 hours.

30 16. Method according to claim 14 to 15, in which the contact temperature used in step b1) is between 15°C and 25°C .

17. Method according to any one of the preceding claims, in which the oxidizing agent is chosen from oxone, potassium permanganate, sodium

percarbonate, peroxides such as hydrogen peroxide, tert-butyl hydroperoxide and m-chloroperoxybenzoic acid.

18. Method according to claim 17, in which the oxidizing agent is hydrogen
5 peroxide.

19. Method according to claim 18, in which the hydrogen peroxide is added in the form of a 35% aqueous solution.

10 20. Method according to any one of the preceding claims, in which the oxidizing agent is used in an amount of 1 to 1.1 molar equivalent.

21. Method according to any one of the preceding claims, in which the reaction temperature in step (ii) is between 28°C and 37°C.

15

22. Method according to one of claims 3 to 21, in which an additional quantity of inorganic acid is added in step (ii).

23. Method according to claim 22, in which the additional quantity of
20 inorganic acid is from 0.02 to 0.3 molar equivalents.

24. Method according to either of claims 22 and 23, in which the contact time in step (ii) is between 10 and 13 hours.

25 25. Method according to any one of the preceding claims, which comprises an additional step (iii) recovering the methyl 2-diphenyl-methylsulfinylacetate obtained.

26. Method according to claim 25, in which step (iii) comprises a distillation
30 of the solvent to dryness.

27. Method according to any one of claims 25 to 26, in which step (iii) comprises a step of direct crystallization.

28. Method according to claim 27, in which the crystallization solvent is chosen from methanol, ethanol, ethyl acetate, isopropyl acetate and toluene.

5 29. Method according to claim 28, in which the crystallization solvent is isopropyl acetate.

30. Method according to any one of the preceding claims, in which the successive steps are carried out in the same reactor without isolation of the
10 intermediate compounds.

31. Method for preparing modafinil comprising preparing MDMSA according to claims 1 to 30.

INTERNATIONAL SEARCH REPORT

PCT/IB2004/000002

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07C317/44 C07C315/02 C07C323/52 C07C319/14

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07C

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

CHEM ABS Data, BEILSTEIN Data, EPO-Internal, PAJ, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US 5 571 825 A (D.H. BOSCHELLI ET AL) 5 November 1996 (1996-11-05) * column 6, lines 64-67 and column 7, lines 4-10; examples 1-2 *	1,17,18, 20,25, 27,31
Y	US 4 964 893 A (L.H. BRANNIGAN ET AL) 23 October 1990 (1990-10-23) * column 21; methods "N", "O" and table I, compounds 45, 54, 61 *	1,17,18, 20,25, 27,31
Y	GB 1 584 462 A (LABORATOIRE L. LAFON) 11 February 1981 (1981-02-11) * example 1a *	1,17,18, 20,25, 27,31

	--- -/-	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

A document defining the general state of the art which is not considered to be of particular relevance

E earlier document but published on or after the international filing date

L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

O document referring to an oral disclosure, use, exhibition or other means

P document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

G document member of the same patent family

Date of the actual completion of the international search

25 May 2004

Date of mailing of the international search report

14/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Rufet, J

INTERNATIONAL SEARCH REPORT

PCT/IB2004/000002

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	N.T. FARINACCI ET AL: J. AM. CHEM. SOC., vol. 59, 1937, pages 2542-2546, XP002234650 * page 2546, column 2, lines 9-14 *	1-3,8,9
A	DATABASE CAPLUS 'Online! CHEMICAL ABSTRACTS SERVICE, COLUMBUS, OHIO, US; retrieved from STN Database accession no. 1996:629856 XP002234652 abstract & JP 08 198843 A (FUJIREBIO KK) 6 August 1996 (1996-08-06)	1,2,14
A	I. SAIKAWA ET AL: CHEM. PHARM. BULL., vol. 33, no. 12, 1985, pages 5534-5538, XP002234651 * page 5535, table II, entry 1d *	1,2,14
A	H. LEHR ET AL: J. MED. CHEM., vol. 6, 1963, pages 136-141, XP001148333 * page 139, experimental, lines 11-25 *	1,2,14

INTERNATIONAL SEARCH REPORT

PCT/IB2004/000002

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5571825	A	05-11-1996	NONE	
US 4964893	A	23-10-1990	CN 87102879 A ,B US 5162537 A US 5321000 A	28-10-1987 10-11-1992 14-06-1994
GB 1584462	A	11-02-1981	BE 865468 A1 CA 1091679 A1 CH 628026 A5 DE 2809625 A1 DK 140878 A ,B, ES 468378 A1 FR 2385693 A1 IE 46566 B1 IT 1094041 B JP 1400453 C JP 53121724 A JP 62009103 B LU 79335 A1 LU 90153 A9 NL 7803432 A ,C US 4177290 A	02-10-1978 16-12-1980 15-02-1982 05-10-1978 01-10-1978 16-12-1978 27-10-1978 27-07-1983 26-07-1985 28-09-1987 24-10-1978 26-02-1987 28-09-1978 16-02-1998 03-10-1978 04-12-1979
JP 08198843	A	06-08-1996	JP 3233806 B2	04-12-2001

THIS PAGE BLANK (USPTO)